

Integrable Dispersive Chains

Maxim V. Pavlov

Novosibirsk State University

28.02.2017

The General Problem. The Korteweg de Vries equation

- The Korteweg de Vries equation is associated with the linear Schrödinger equation

$$\psi_{xx} = (\lambda + u)\psi.$$

The function $\psi(x, t, \lambda)$ satisfies the pair of *linear* equations in partial derivatives

$$\psi_{xx} = u\psi, \quad \psi_t = a\psi_x - \frac{1}{2}a_x\psi.$$

Then the compatibility condition $(\psi_{xx})_t = (\psi_t)_{xx}$ yields the relationship

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x \right) a$$

between functions $u(x, t, \lambda)$ and $a(x, t, \lambda)$.

The General Problem. The Korteweg de Vries equation

- The Korteweg de Vries equation is associated with the linear Schrödinger equation

$$\psi_{xx} = (\lambda + u)\psi.$$

The function $\psi(x, t, \lambda)$ satisfies the pair of *linear* equations in partial derivatives

$$\psi_{xx} = u\psi, \quad \psi_t = a\psi_x - \frac{1}{2}a_x\psi.$$

Then the compatibility condition $(\psi_{xx})_t = (\psi_t)_{xx}$ yields the relationship

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x \right) a$$

between functions $u(x, t, \lambda)$ and $a(x, t, \lambda)$.

- If we choose the linear dependences $u(x, t, \lambda) = \lambda + u^1(x, t)$ and $a(x, t, \lambda) = \lambda + a_1(x, t)$, we obtain nothing but the famous Korteweg de Vries equation

$$u_t^1 = \frac{1}{4}u_{xxx}^1 - \frac{3}{2}u^1u_x^1,$$

- Again we consider the relationship

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x \right) a$$

between functions $u(x, t, \lambda)$ and $a(x, t, \lambda)$.

The General Problem. The Kaup–Boussinesq system

- Again we consider the relationship

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x \right) a$$

between functions $u(x, t, \lambda)$ and $a(x, t, \lambda)$.

- If we choose the quadratic dependence $u(x, t, \lambda) = \lambda^2 + \lambda u^1(x, t) + u^2(x, t)$ and again the linear dependence $a(x, t, \lambda) = \lambda + a_1(x, t)$, we obtain nothing but the well-known Kaup–Boussinesq system

$$u_t^1 = u_x^2 - \frac{3}{2}u^1 u_x^1, \quad u_t^2 = \frac{1}{4}u_{xxx}^1 - u^2 u_x^1 - \frac{1}{2}u^1 u_x^2.$$

The General Problem. The Antonowicz–Fordy Construction

- Again we consider the relationship

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x \right) a$$

between functions $u(x, t, \lambda)$ and $a(x, t, \lambda)$.

The General Problem. The Antonowicz–Fordy Construction

- Again we consider the relationship

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x \right) a$$

between functions $u(x, t, \lambda)$ and $a(x, t, \lambda)$.

- Multi-component *rational* (with respect to the spectral parameter λ) generalization (ϵ_k are arbitrary parameters)

$$u(x, t, \lambda) = \frac{\lambda^M u^0(x, t) + \lambda^{M-1} u^1(x, t) + \dots + u^M(x, t)}{\epsilon_M \lambda^M + \epsilon_{M-1} \lambda^{M-1} + \dots + \epsilon_0}.$$

Integrable Dispersive Chains

- Now we consider ($M = 1, 2, \dots$)

$$u(x, \mathbf{t}, \lambda) = \lambda^M \left(1 + \frac{u^1(x, \mathbf{t})}{\lambda} + \frac{u^2(x, \mathbf{t})}{\lambda^2} + \frac{u^3(x, \mathbf{t})}{\lambda^3} + \dots \right),$$

where u^k are infinitely many unknown functions.

Integrable Dispersive Chains

- Now we consider ($M = 1, 2, \dots$)

$$u(x, \mathbf{t}, \lambda) = \lambda^M \left(1 + \frac{u^1(x, \mathbf{t})}{\lambda} + \frac{u^2(x, \mathbf{t})}{\lambda^2} + \frac{u^3(x, \mathbf{t})}{\lambda^3} + \dots \right),$$

where u^k are infinitely many unknown functions.

- The substitution and the linear dependence $a^{(1)} = \lambda + a_1(x, \mathbf{t})$ into

$$u_t = \left(-\frac{1}{2} \partial_x^3 + 2u \partial_x + u_x \right) a$$

yields M th integrable dispersive chain

$$u_t^k = u_x^{k+1} - \frac{1}{2} u^1 u_x^k - u^k u_x^1 + \frac{1}{4} \delta_M^k u_{xxx}^1, \quad k = 1, 2, \dots,$$

where δ_M^k is the Kronecker delta and

$$a_1 = -\frac{1}{2} u^1.$$

Higher Commuting Flows

- Higher commuting flows of the Korteweg de Vries hierarchy are determined by the linear spectral system

$$\psi_{xx} = (\lambda + u^1)\psi, \quad \psi_{t^k} = a^{(k)}\psi_x - \frac{1}{2}a_x^{(k)}\psi,$$

where

$$a^{(k)} = \lambda^k + \sum_{m=1}^k a_m \lambda^{k-m},$$

and functions a_m and u^1 depend on the “space” variable x and infinitely many extra “time” variables t^k (obviously, $t \equiv t^1$).

Higher Commuting Flows

- Higher commuting flows of the Korteweg de Vries hierarchy are determined by the linear spectral system

$$\psi_{xx} = (\lambda + u^1)\psi, \quad \psi_{t^k} = a^{(k)}\psi_x - \frac{1}{2}a_x^{(k)}\psi,$$

where

$$a^{(k)} = \lambda^k + \sum_{m=1}^k a_m \lambda^{k-m},$$

and functions a_m and u^1 depend on the “space” variable x and infinitely many extra “time” variables t^k (obviously, $t \equiv t^1$).

- Substitution

$$u(x, \mathbf{t}, \lambda) = \lambda^M \left(1 + \frac{u^1(x, \mathbf{t})}{\lambda} + \frac{u^2(x, \mathbf{t})}{\lambda^2} + \frac{u^3(x, \mathbf{t})}{\lambda^3} + \dots \right),$$

into

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x \right) a$$

Higher Commuting Flows

leads to higher commuting flows (here we define $a_0 = 1$)

$$u_{t^s}^k = \sum_{m=0}^s \left(u^{k+m} \partial_x + \partial_x u^{k+m} - \frac{1}{2} \delta_M^{k+m} \partial_x^3 \right) a_{s-m}, \quad s = 1, 2, \dots,$$

Higher Commuting Flows

leads to higher commuting flows (here we define $a_0 = 1$)

$$u_{t^s}^k = \sum_{m=0}^s \left(u^{k+m} \partial_x + \partial_x u^{k+m} - \frac{1}{2} \delta_M^{k+m} \partial_x^3 \right) a_{s-m}, \quad s = 1, 2, \dots,$$

where all coefficients a_m can be found iteratively from the linear system (here we define $u^0 = 1$ and $u^{-m} = 0$ for all $m = 1, 2, \dots$)

$$\sum_{m=0}^s \left(u^{m-k} \partial_x + \partial_x u^{m-k} - \frac{1}{2} \delta_M^{m-k} \partial_x^3 \right) a_{s-m} = 0, \quad k = 0, 1, \dots, s-1.$$

For instance,

$$a_1 = -\frac{1}{2} u^1, \quad a_2 = -\frac{1}{2} u^2 + \frac{3}{8} (u^1)^2 - \frac{1}{8} \delta_M^1 u_{xx}^1, \quad a_3 = -\frac{1}{2} u^3 + \frac{3}{4} u^1 u^2 - \frac{5}{16} (u^1)^3 + \frac{1}{32} \delta_M^1 (10 u^1 u_{xx}^1 + 5 (u_x^1)^2 - u_{xxxx}^1 - 4 u_{xx}^2) - \frac{1}{8} \delta_M^2 u_{xx}^1, \dots$$

Higher Commuting Flows

Thus all higher commuting flows are written also in an evolution form. For instance, the first commuting flow to

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1 + \frac{1}{4}\delta_M^k u_{xxx}^1, \quad k = 1, 2, \dots,$$

is (here we identify $y \equiv t^2$)

$$\begin{aligned} u_y^k &= u_x^{k+2} - \frac{1}{2}u^1 u_x^{k+1} + \left(-\frac{1}{2}u^2 + \frac{3}{8}(u^1)^2 - \frac{1}{8}\delta_M^1 u_{xx}^1 \right) u_x^k - u^{k+1} u_x^1 \\ &\quad + u^k \left(-u_x^2 + \frac{3}{2}u^1 u_x^1 - \frac{1}{4}\delta_M^1 u_{xxx}^1 \right) + \frac{1}{4}\delta_M^{k+1} u_{xxx}^1 \\ &\quad + \frac{1}{4}\delta_M^k \left(u_{xxx}^2 - \frac{3}{4}[(u^1)^2]_{xxx} + \frac{1}{4}\delta_M^1 u_{xxxxx}^1 \right). \end{aligned}$$

Local Hamiltonian Structures

- A hierarchy of integrable dispersive chains

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1 + \frac{1}{4}\delta_M^k u_{xxx}^1, \quad k = 1, 2, \dots,$$

possesses *infinitely* many local Hamiltonian structures:

$$u_{t^s}^k = \sum_{m=1}^{s+1} \left(u^{k+m-1} \partial_x + \partial_x u^{k+m-1} - \frac{1}{2} \delta_M^{k+m-1} \partial_x^3 \right) \frac{\delta \mathbf{H}_{s+1}}{\delta u^m};$$

$$u_{t^s}^1 = -2 \partial_x \frac{\delta \mathbf{H}_{s+2}}{\delta u^1},$$

$$u_{t^s}^k = \sum_{m=2}^{s+2} \left(u^{k+m-2} \partial_x + \partial_x u^{k+m-2} - \frac{1}{2} \delta_M^{k+m-2} \partial_x^3 \right) \frac{\delta \mathbf{H}_{s+2}}{\delta u^m};$$

$$u_{t^s}^1 = -2 \partial_x \frac{\delta \mathbf{H}_{s+3}}{\delta u^2}, \quad u_{t^s}^2 = -2 \partial_x \frac{\delta \mathbf{H}_{s+3}}{\delta u^1} - \left(u^1 \partial_x + \partial_x u^1 - \frac{1}{2} \delta_M^1 \partial_x^3 \right) \frac{\delta \mathbf{H}_{s+3}}{\delta u^2},$$

$$u_{t^s}^k = \sum_{m=3}^{s+3} \left(u^{k+m-3} \partial_x + \partial_x u^{k+m-3} - \frac{1}{2} \delta_M^{k+m-3} \partial_x^3 \right) \frac{\delta \mathbf{H}_{s+3}}{\delta u^m}.$$

- All higher local conservation laws can be found from the observation

$$a_m = \frac{\delta \mathbf{H}_{m+s}}{\delta u^s}, \quad m = 0, 1, \dots; \quad s = 1, 2, \dots$$

In such a case all Hamiltonians can be found from above variation derivatives, for instance

$$\mathbf{H}_1 = \int u^1 dx, \quad \mathbf{H}_2 = \int \left(u^2 - \frac{1}{4}(u^1)^2 \right) dx,$$

$$\mathbf{H}_3 = \int \left(u^3 - \frac{1}{2}u^1 u^2 + \frac{1}{8}(u^1)^3 + \frac{1}{16}\delta_M^1(u_x^1)^2 \right) dx,$$

$$\begin{aligned} \mathbf{H}_4 = & \int \left(u^4 - \frac{1}{2}u^1 u^3 - \frac{1}{4}(u^2)^2 + \frac{3}{8}(u^1)^2 u^2 - \frac{5}{64}(u^1)^4 \right. \\ & \left. + \frac{1}{32}\delta_M^1 \left(-5u^1 (u_x^1)^2 - \frac{1}{2}(u_{xx}^1)^2 + 4u_x^1 u_x^2 \right) + \frac{1}{16}\delta_M^2 (u_x^1)^2 \right) dx, \dots \end{aligned}$$

Elementary Reductions

Obviously for any natural number $N \geq M$ the reduction $u^{N+1} = 0$ of M th dispersive chain

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1 + \frac{1}{4}\delta_M^k u_{xxx}^1, \quad k = 1, 2, \dots,$$

leads to N component integrable dispersive systems:

Elementary Reductions

Obviously for any natural number $N \geq M$ the reduction $u^{N+1} = 0$ of M th dispersive chain

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1 + \frac{1}{4}\delta_M^k u_{xxx}^1, \quad k = 1, 2, \dots,$$

leads to N component integrable dispersive systems:

1. $N = M = 1$, the Korteweg de Vries equation;

Elementary Reductions

Obviously for any natural number $N \geq M$ the reduction $u^{N+1} = 0$ of M th dispersive chain

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1 + \frac{1}{4}\delta_M^k u_{xxx}^1, \quad k = 1, 2, \dots,$$

leads to N component integrable dispersive systems:

1. $N = M = 1$, the Korteweg de Vries equation;
2. $N = 2, M = 1$, the Ito system

$$u_t^1 = u_x^2 - \frac{3}{2}u^1 u_x^1 + \frac{1}{4}u_{xxx}^1, \quad u_t^2 = -\frac{1}{2}u^1 u_x^2 - u^2 u_x^1;$$

Elementary Reductions

Obviously for any natural number $N \geq M$ the reduction $u^{N+1} = 0$ of M th dispersive chain

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1 + \frac{1}{4}\delta_M^k u_{xxx}^1, \quad k = 1, 2, \dots,$$

leads to N component integrable dispersive systems:

1. $N = M = 1$, the Korteweg de Vries equation;
2. $N = 2, M = 1$, the Ito system

$$u_t^1 = u_x^2 - \frac{3}{2}u^1 u_x^1 + \frac{1}{4}u_{xxx}^1, \quad u_t^2 = -\frac{1}{2}u^1 u_x^2 - u^2 u_x^1;$$

3. $N > 2, M = 1$,

$$u_t^1 = u_x^2 - \frac{3}{2}u^1 u_x^1 + \frac{1}{4}u_{xxx}^1,$$

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1, \quad k = 2, \dots, N-1,$$

$$u_t^N = -\frac{1}{2}u^1 u_x^N - u^N u_x^1;$$

4. $N = M > 1$, (if $N = M = 2$, this is the Kaup–Boussinesq equation)

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1, \quad k = 1, 2, \dots, N-1,$$

$$u_t^N = -\frac{1}{2}u^1 u_x^N - u^N u_x^1 + \frac{1}{4}u_{xxx}^1;$$

Elementary Reductions

4. $N = M > 1$, (if $N = M = 2$, this is the Kaup–Boussinesq equation)

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1, \quad k = 1, 2, \dots, N-1,$$

$$u_t^N = -\frac{1}{2}u^1 u_x^N - u^N u_x^1 + \frac{1}{4}u_{xxx}^1;$$

5. $N = M + 1, M > 1$,

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1, \quad k = 1, 2, \dots, N-2,$$

$$u_t^{N-1} = u_x^N - \frac{1}{2}u^1 u_x^{N-1} - u^{N-1} u_x^1 + \frac{1}{4}u_{xxx}^1,$$

$$u_t^N = -\frac{1}{2}u^1 u_x^N - u^N u_x^1.$$

6. $N > M + 1, M > 1,$

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1, \quad k = 1, \dots, M,$$

$$u_t^M = u_x^{M+1} - \frac{1}{2}u^1 u_x^M - u^M u_x^1 + \frac{1}{4}u_{xxx}^1,$$

$$u_t^k = u_x^{k+1} - \frac{1}{2}u^1 u_x^k - u^k u_x^1, \quad k = M + 1, \dots, N - 1,$$

$$u_t^N = -\frac{1}{2}u^1 u_x^N - u^N u_x^1.$$

Rational Factorised Constraints

- Now we consider more complicated N component reductions
($M = 1, 2, \dots, K = 0, 1, \dots$)

$$u(x, \mathbf{t}, \lambda) = \frac{\lambda^{M+K} + \lambda^{M+K-1} v_{M+K-1}(x, \mathbf{t}) + \dots + \lambda v_1(x, \mathbf{t}) + v_0(x, \mathbf{t})}{\lambda^K + \lambda^{K-1} w_{K-1} + \dots + \lambda w_1 + w_0}$$

where w_k are arbitrary constants.

Rational Factorised Constraints

- Now we consider more complicated N component reductions ($M = 1, 2, \dots, K = 0, 1, \dots$)

$$u(x, \mathbf{t}, \lambda) = \frac{\lambda^{M+K} + \lambda^{M+K-1} v_{M+K-1}(x, \mathbf{t}) + \dots + \lambda v_1(x, \mathbf{t}) + v_0(x, \mathbf{t})}{\lambda^K + \lambda^{K-1} w_{K-1} + \dots + \lambda w_1 + w_0}$$

where w_k are arbitrary constants.

- Suppose for simplicity that all roots of these two polynomials are pairwise distinct. Then the substitution $a^{(1)} = \lambda + a_1(x, \mathbf{t})$ and (here r^k are arbitrary constants)

$$u(x, \mathbf{t}, \lambda) = \frac{\prod_{m=1}^{M+K} (\lambda - s^m(x, \mathbf{t}))}{\prod_{k=1}^K (\lambda - r^k)}$$

into

$$u_t = \left(-\frac{1}{2} \partial_x^3 + 2u \partial_x + u_x \right) a$$

Rational Factorised Constraints

- yields multi-component integrable dispersive systems:

$$s_t^i = (s^i + a_1) s_x^i + \frac{1}{2} \frac{\prod_{k=1}^K (s^i - r^k)}{\prod_{m \neq i} (s^i - s^m)} a_{1,xxx},$$

where

$$a_1 = \frac{1}{2} \left(\sum_{m=1}^{M+K} s^m - \sum_{k=1}^K r^k \right).$$

Rational Factorised Constraints

- yields multi-component integrable dispersive systems:

$$s_t^i = (s^i + a_1)s_x^i + \frac{1}{2} \frac{\prod_{k=1}^K (s^i - r^k)}{\prod_{m \neq i} (s^i - s^m)} a_{1,xxx},$$

where

$$a_1 = \frac{1}{2} \left(\sum_{m=1}^{M+K} s^m - \sum_{k=1}^K r^k \right).$$

- For instance, the Kaup–Boussinesq system becomes

$$s_t^1 = \frac{1}{2} (3s^1 + s^2)s_x^1 + \frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}, \quad s_t^2 = \frac{1}{2} (s^1 + 3s^2)s_x^2 - \frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)};$$

the Ito system takes the form

$$s_t^1 = \frac{1}{2} (3s^1 + s^2)s_x^1 + \frac{s^1(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}, \quad s_t^2 = \frac{1}{2} (s^1 + 3s^2)s_x^2 - \frac{s^2(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}.$$

Three Dimensional Linearly Degenerate Quasilinear Equations

- The three dimensional quasilinear system

$$a_{1,t} = a_{2,x}, \quad a_1 a_{2,x} + a_{1,y} = a_2 a_{1,x} + a_{2,t},$$

is determined by the compatibility conditions

$$p_t = [(\lambda + a_1)p]_x, \quad p_y = [(\lambda^2 + a_1\lambda + a_2)p]_x,$$

Three Dimensional Linearly Degenerate Quasilinear Equations

- The three dimensional quasilinear system

$$a_{1,t} = a_{2,x}, \quad a_1 a_{2,x} + a_{1,y} = a_2 a_{1,x} + a_{2,t},$$

is determined by the compatibility conditions

$$p_t = [(\lambda + a_1)p]_x, \quad p_y = [(\lambda^2 + a_1\lambda + a_2)p]_x,$$

where $p = 1/\varphi$. Here $\varphi = \psi\psi^+$, where ψ and ψ^+ are two linearly independent solutions of

$$\psi_{xx} = u\psi,$$

$$\psi_t = (\lambda + a_1)\psi_x - \frac{1}{2}a_{1,x}\psi, \quad \psi_y = (\lambda^2 + a_1\lambda + a_2)\psi_x - \frac{1}{2}(\lambda a_{1,x} + a_{2,x})\psi.$$

- **Statement:** *Three dimensional quasilinear system*

$$a_{1,t} = a_{2,x}, \quad a_1 a_{2,x} + a_{1,y} = a_2 a_{1,x} + a_{2,t}$$

possesses infinitely many finite component differential constraints $a_1(\mathbf{u}, \mathbf{u}_x, \dots)$, $a_2(\mathbf{u}, \mathbf{u}_x, \dots)$, where field variables u^k are solutions of dispersive integrable systems determined by linear spectral problem

$$\psi_{xx} = u\psi,$$

$$\psi_t = (\lambda + a_1)\psi_x - \frac{1}{2}a_{1,x}\psi, \quad \psi_y = (\lambda^2 + a_1\lambda + a_2)\psi_x - \frac{1}{2}(\lambda a_{1,x} + a_{2,x})\psi,$$

where

$$u(x, t, y, \lambda) = \frac{\prod_{m=1}^{M+K} (\lambda - s^m(x, t, y))}{\prod_{k=1}^K (\lambda - r^k)}.$$

Dispersive Reductions

Differential constraints

$$a_1 = \frac{1}{2} \left(\sum_{m=1}^{M+K} s^m - \sum_{k=1}^K r^k \right), \quad a_2 = \frac{1}{4} \sum_{m=1}^{M+K} (s^m)^2 - \frac{1}{4} \sum_{k=1}^K (r^k)^2 + \frac{1}{2} a_1^2 + \frac{1}{4} \delta_M^1 a_{1,xx}$$

and integrable commuting flows

$$s_t^i = (s^i + a_1) s_x^i + \frac{1}{2} \frac{\prod_{k=1}^K (s^i - r^k)}{\prod_{m \neq i} (s^i - s^m)} a_{1,xxx},$$

$$s_y^i = (a_2 + a_1 s^i + (s^i)^2) s_x^i + \frac{1}{2} \frac{\prod_{k=1}^K (s^i - r^k)}{\prod_{m \neq i} (s^i - s^m)} (s^i a_{1,xxx} + a_{2,xxx}).$$

Differential Constraints. The Korteweg de Vries equation

Differential constraints are

$$a_1 = -\frac{1}{2}u^1, \quad a_2 = \frac{3}{8}(u^1)^2 - \frac{1}{8}u_{xx}^1.$$

Substitution into the three dimensional quasilinear system

$$a_{1,t} = a_{2,x}, \quad a_1 a_{2,x} + a_{1,y} = a_2 a_{1,x} + a_{2,t}$$

leads to an identity, if $u^1(x, t, y)$ is an arbitrary solution of the Korteweg de Vries equation

$$u_t^1 = \left(\frac{1}{4}u_{xx}^1 - \frac{3}{4}(u^1)^2 \right)_x$$

and into its first commuting flow

$$u_y^1 = \left(\frac{5}{8}(u^1)^3 - \frac{5}{16}(u_x^1)^2 - \frac{5}{8}u^1 u_{xx}^1 + \frac{1}{16}u_{xxxx}^1 \right)_x.$$